Surname	Other na	nmes
Pearson Edexcel GCE	Centre Number	Candidate Number
Mechani Advanced/Advan		
Friday 17 June 2016 – A Time: 1 hour 30 minut		Paper Reference 6678/01

Candidates may use any calculator allowed by the regulations of the Joint Council for Qualifications. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B). Coloured pencils and highlighter pens must not be used.
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer all questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided
 there may be more space than you need.
- You should show sufficient working to make your methods clear. Answers without working may not gain full credit.
- Whenever a numerical value of g is required, take g = 9.8 m s⁻², and give your answer to either two significant figures or three significant figures.
- When a calculator is used, the answer should be given to an appropriate degree of accuracy.

Information

- The total mark for this paper is 75.
- The marks for **each** question are shown in brackets
 - use this as a guide as to how much time to spend on each question.

Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.

PEARSON

Turn over ▶

P46710A

©2016 Pearson Education Ltd.

1.	v m	particle P moves along a straight line. The speed of P at time t seconds t s ⁻¹ , where t = (t = t = t = t and t = t	
	1.1111	u	
	(a)	the acceleration of P when $t = 3$	(8)
	(b)	the distance travelled by P in the third second of the motion.	(5)

	Leave
	blank
	Dialik
Question 1 continued	
	1



	Leave blank
Question 1 continued	

Q1 (Total 13 marks)	Question 1 continued	Leave
	Question I continued	
(Total 13 marks)		Q1
	(Total 13 marks)	



- 2. A car of mass 800kg is moving on a straight road which is inclined at an angle θ to the horizontal, where $\sin \theta = \frac{1}{20}$. The resistance to the motion of the car from non-gravitational forces is modelled as a constant force of magnitude R newtons. When the car is moving up the road at a constant speed of $12.5 \,\mathrm{m\,s^{-1}}$, the engine of the car is working at a constant rate of 3P watts. When the car is moving down the road at a constant speed of $12.5 \,\mathrm{m\,s^{-1}}$, the engine of the car is working at a constant rate of P watts.
 - (a) Find
 - (i) the value of P,
 - (ii) the value of R.

(6)

When the car is moving up the road at $12.5\,\mathrm{m\,s^{-1}}$ the engine is switched off and the car comes to rest, without braking, in a distance d metres. The resistance to the motion of the car from non-gravitational forces is still modelled as a constant force of magnitude R newtons.

(b) Use the work-energy principle to find the value of d .	

estion 2 continued	



	Leave
	blank
Question 2 continued	

Question 2 continued	Leave
Question 2 continued	
	Q2
(Total 10 marks)	



3.	A particle of mass 0.6kg is moving with constant velocity $(c\mathbf{i} + 2c\mathbf{j}) \mathrm{ms^{-1}}$, where c is a positive constant. The particle receives an impulse of magnitude $2\sqrt{10}\mathrm{Ns}$. Immediately after receiving the impulse the particle has velocity $(2c\mathbf{i} - c\mathbf{j}) \mathrm{ms^{-1}}$.	
	immediately after receiving the impulse the particle has velocity (201 – cj) his .	
	Find the value of c .	
	(6)	

Question 3 continued		Leave
		Q3
	(Total 6 marks)	



4.




Figure 1

The uniform lamina OBC is one quarter of a circular disc with centre O and radius Am. The points A and A0, on A0 and A1 are A2 and A3 and A4 and A5. The uniform lamina A4 and A5 shown shaded in Figure 1, is formed by removing the triangle A4 from A5.

Given that the centre of mass of one quarter of a uniform circular disc of radius r is at a distance $\frac{4\sqrt{2}}{3\pi}r$ from the centre of the disc,

(a) find the distance of the centre of mass of the lamina ABCD from AD.

(5)

The lamina is freely suspended from D and hangs in equilibrium.

(b) Find, to the nearest degree, the angle between DC and the downward vertical. (4)



	L
Question 4 continued	0
(



Question 4 continued		

	blank
Question 4 continued	
	04
	Q4
(Total 9 marks)	



5.




Figure 2

A non-uniform rod AB, of mass 5kg and length 4m, rests with one end A on rough horizontal ground. The centre of mass of the rod is d metres from A. The rod is held in limiting equilibrium at an angle θ to the horizontal by a force \mathbf{P} , which acts in a direction perpendicular to the rod at B, as shown in Figure 2. The line of action of \mathbf{P} lies in the same vertical plane as the rod.

- (a) Find, in terms of d, g and θ ,
 - (i) the magnitude of the vertical component of the force exerted on the rod by the ground,
 - (ii) the magnitude of the friction force acting on the rod at A.

(8)

Given that $\tan \theta = \frac{5}{12}$ and that the coefficient of friction between the rod and the ground is $\frac{1}{2}$,

(b) find the value of d.

(4)

	Leave blank
Question 5 continued	



Question 5 continued	
Question 6 continue	

	Leave blank
Question 5 continued	
	_
	-
	-
	-
	_
	_
	_
	_
	-
	-
	-
	_
	_
	_
	_
	_
	-
	-
	-
	-
	_
	_
	_
	_
	-
	-
	-
	-
	_
	_
	Q5
(Total 12 marks	,
(10tal 12 marks	



Leave

6.	ΓIn	this question, i is a horizontal unit vector and j is an upward vertical unit vector.]
0.	A p	article P is projected from a fixed origin O with velocity $(3\mathbf{i} + 4\mathbf{j}) \mathrm{m \ s^{-1}}$. The particle ves freely under gravity and passes through the point A with position vector $\lambda(\mathbf{i} - \mathbf{j}) \mathrm{m}$, ere λ is a positive constant.
	(a)	Find the value of λ .
		(6)
	(b)	Find
		(i) the speed of P at the instant when it passes through A ,
		(ii) the direction of motion of P at the instant when it passes through A . (7)



uestion 6 continued	



Question 6 continued	
Question o commune	
	_
	_
	_
	_
	_

	Leave
Question 6 continued	blank
Question o continued	
	—
	—
	—
	—
	—
	_
	—
	Q6
(Total 13 mar	ks)



- 7. Two particles A and B, of mass 2m and 3m respectively, are initially at rest on a smooth horizontal surface. Particle A is projected with speed 3u towards B. Particle A collides directly with particle B. The coefficient of restitution between A and B is $\frac{3}{4}$
 - (a) Find
 - (i) the speed of A immediately after the collision,
 - (ii) the speed of B immediately after the collision.

(7)

After the collision B hits a fixed smooth vertical wall and rebounds. The wall is perpendicular to the direction of motion of B. The coefficient of restitution between B and the wall is e. The magnitude of the impulse received by B when it hits the wall is $\frac{27}{4}$ mu.

(b) Find the value of e.

(3)

(c) Determine whether there is a further collision between *A* and *B* after *B* rebounds from the wall.

(2)

uestion 7 continued	



Question 7 continued	

	Leave
	blank
Question 7 continued	



	(Total 12 marks) TOTAL FOR PAPER: 75 MARKS	