Surname	Other na	ames
Pearson Edexcel GCE	Centre Number	Candidate Number
Mechani Advanced/Advan		
Tuesday 21 June 2016 – Time: 1 hour 30 minut	•	Paper Reference 6681/01

Candidates may use any calculator allowed by the regulations of the Joint Council for Qualifications. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B). Coloured pencils and highlighter pens must not be used.
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided
 there may be more space than you need.
- You should show sufficient working to make your methods clear. Answers without working may not gain full credit.
- Whenever a numerical value of g is required, take $g = 9.8 \text{ m s}^{-2}$, and give your answer to either two significant figures or three significant figures.
- When a calculator is used, the answer should be given to an appropriate degree of accuracy.

Information

- The total mark for this paper is 75.
- The marks for **each** question are shown in brackets
 - use this as a guide as to how much time to spend on each question.

Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.

PEARSON

Turn over ▶



Leave blank

1.	[In this question, \mathbf{i} and \mathbf{j} are perpendicular unit vectors in a horizontal plane.]
	A bead P of mass 0.4 kg is threaded on a smooth straight horizontal wire. The wire lies along the line with vector equation $\mathbf{r} = (\mathbf{i} + 2\mathbf{j}) + \lambda(-2\mathbf{i} + 3\mathbf{j})$. The bead is initially at rest at the point A with position vector $(-\mathbf{i} + 5\mathbf{j})$ m. A constant horizontal force $(0.5\mathbf{i} + \mathbf{j})$ N acts on P and moves it along the wire to the point B . At B the speed of P is 5 m s ⁻¹ .
	Find the position vector of <i>B</i> .
	(7)



Question 1 continued	Leave blank
	Q1
(Total 7 marks)	



2. A particle *P* is moving in a plane. At time *t* seconds the position vector of *P* is **r** metres and the velocity of *P* is **v** m s⁻¹. When $t = \frac{\pi}{2}$, *P* is instantaneously at rest at the point with position vector $(\mathbf{i} - \mathbf{j})$ m.

Given that \mathbf{r} satisfies the differential equation

$$\frac{\mathrm{d}^2\mathbf{r}}{\mathrm{d}t^2} + 4\mathbf{r} = (3\sin t)\,\mathbf{i}$$

find **v** in terms of *t*.

(13)

Question 2 continued	blank
	02
	Q2
(Total 13 marks)	



Three forces \mathbf{F}_1 , \mathbf{F}_2 and \mathbf{F}_3 act on a rigid body at the points with position vectors \mathbf{r}_1 , \mathbf{r}_2 and \mathbf{r}_3 respectively, where

$$\mathbf{F}_{1} = (2\mathbf{j} - \mathbf{k}) \text{ N}$$
 $\mathbf{F}_{2} = (\mathbf{i} + \mathbf{k}) \text{ N}$ $\mathbf{F}_{3} = (\mathbf{i} + \mathbf{j}) \text{ N}$

$$\mathbf{F}_2 = (\mathbf{i} + \mathbf{k}) \, \mathbf{N}$$

$$\mathbf{F}_3 = (\mathbf{i} + \mathbf{j}) \, \mathbf{N}$$

$$\mathbf{r}_{i} = (4\mathbf{i} - \mathbf{k}) \, \mathbf{n}$$

$$\mathbf{r}_2 = (2\mathbf{i} + \mathbf{k}) \, \mathbf{n}$$

$$\mathbf{r}_1 = (4\mathbf{j} - \mathbf{k}) \text{ m}$$
 $\mathbf{r}_2 = (2\mathbf{i} + \mathbf{k}) \text{ m}$ $\mathbf{r}_3 = (3\mathbf{i} + \mathbf{j} + \mathbf{k}) \text{ m}$

The system of the three forces is equivalent to a single force R acting through the point with position vector $(\mathbf{i} - \mathbf{j} + \mathbf{k})$ m, together with a couple of moment \mathbf{G} .

(a) Find R.

(2)

(h) Find G

(9)

(0)	1	ma	u.

6

	blank
Question 3 continued	
	Q3
(Total 11 marks)	



Leave blank

height h and mass M , about a diameter of one end.	(10
	(10

uestion 4 continued	




Leave blank

Question 4 continued		

Question 4 continued	Leave
	Q4
(Total 10 marks)	




5.



A uniform piece of wire ABC, of mass 2m and length 4a, is bent into two straight equal portions, AB and BC, which are at right angles to each other, as shown in Figure 1. The wire rotates freely in a vertical plane about a fixed smooth horizontal axis L which passes through A and is perpendicular to the plane of the wire.

(a) Show that the moment of inertia of the wire about L is $\frac{20ma^2}{3}$





estion 5 continued	
	_



Leave blank

Question 5 continued	

	blank
Question 5 continued	
	Q5
(Total 11 marks)	
(10tai 11 marks)	



(6)

- 6. A firework rocket, excluding its fuel, has mass m_0 kg. The rocket moves vertically upwards by ejecting burnt fuel vertically downwards with constant speed u m s⁻¹, u > 24.5, relative to the rocket. The rocket starts from rest on the ground at time t = 0. At time t seconds, $t \le 2$, the speed of the rocket is v m s⁻¹ and the mass of the rocket including its fuel is $m_0(5-2t)$ kg. It is assumed that air resistance is negligible and the acceleration due to gravity is constant.
 - (a) Show that, for $t \leq 2$

$$\frac{\mathrm{d}v}{\mathrm{d}t} = \frac{2u}{5 - 2t} - 9.8$$

(b)	Find the speed of the rocket at the instant when all of its fuel has been burnt.	
		(6)



	Leave
	blank
Question 6 continued	
	1



	Leave blank
Question 6 continued	

	Leave
Question 6 continued	
	Q6
(Total 12 marks)	



- 7. A uniform square lamina PQRS, of mass m and side 2a, is free to rotate about a fixed smooth horizontal axis which passes through P and Q. The lamina hangs at rest in a vertical plane with SR below PQ and is given a horizontal impulse of magnitude J at the midpoint of SR. The impulse is perpendicular to SR.
 - (a) Find the initial angular speed of the lamina.

(3)

(b) Find the magnitude of the angular deceleration of the lamina at the instant when the lamina has turned through $\frac{\pi}{6}$ radians.

(3)

(c) Find the magnitude of the component of the force exerted on the lamina by the axis, in a direction perpendicular to the lamina, at the instant when the lamina has turned through $\frac{\pi}{6}$ radians.

(5)

uestion 7 continued	



Leave blank

Question 7 continued	

uestion 7 continued	



	(Total 11 marks)
	TOTAL FOR PAPER: 75 MARKS
T	END