



Rewarding Learning

ADVANCED SUBSIDIARY (AS)

General Certificate of Education

2016

Mathematics

Assessment Unit M1

assessing

Module M1: Mechanics 1



AMM11

[AMM11]

WEDNESDAY 25 MAY, MORNING

TIME

1 hour 30 minutes.

INSTRUCTIONS TO CANDIDATES

Write your Centre Number and Candidate Number on the Answer Booklet provided.

Answer **all seven** questions.

Show clearly the full development of your answers.

Answers should be given to three significant figures unless otherwise stated.

You are permitted to use a graphic or scientific calculator in this paper.

INFORMATION FOR CANDIDATES

The total mark for this paper is 75

Figures in brackets printed down the right-hand side of pages indicate the marks awarded to each question or part question.

Answers should include diagrams where appropriate and marks may be awarded for them.

Take $g = 9.8 \text{ m s}^{-2}$, unless specified otherwise.

A copy of the **Mathematical Formulae and Tables booklet** is provided.

Answer all seven questions.

Show clearly the full development of your answers.

Answers should be given to three significant figures unless otherwise stated.

- 1** A stone is thrown vertically downwards from the top of a cliff at a speed of 5 m s^{-1} .
The top of the cliff is 250 m above horizontal ground.
If air resistance is ignored,

(i) find the speed of the stone as it hits the ground, [2]

(ii) find the time taken for the stone to fall to the ground. [2]

The mass of the stone is 0.2 kg.

If air resistance is included as a constant force of magnitude 0.8 N,

(iii) find the acceleration of the stone. [3]

- 2 **Fig. 1** below shows the displacement-time graph of a cyclist's journey along a straight horizontal road.




Fig. 1

Five points O, A, B, C and D are marked on the graph above corresponding to the cyclist's journey at 0, 120, 180, 270 and 312 seconds respectively.

- (i) Find the velocity of the cyclist as he travels from O to A. [2]

The cyclist maintains a constant velocity as he travels from B to D.

- (ii) Find the displacement of D from O. [4]

- (iii) Find the cyclist's average speed for the complete journey. [3]

- 3 **Fig. 2** below shows a light fitting of mass 2 kg suspended by electric cables AF and FB. The ends A and B are fixed to a horizontal ceiling and a vertical wall respectively. The system is in equilibrium and lies in a vertical plane.




Fig. 2

The cable AF is inclined at an angle of θ° to the vertical and the tension in this cable is T N. The cable FB is inclined at an angle of 20° to the horizontal and the tension in this cable is 10 N.

Model the light fitting as a particle.

- (i) State one modelling assumption you will make about the cables. [1]
- (ii) Draw a diagram showing the external forces acting on the light fitting. [2]
- (iii) Find T and θ . [8]

- 4 A particle P moves in a straight line so that its velocity $v \text{ m s}^{-1}$ at time t seconds, is given by

$$v = 0.6t^2 + kt + 12$$

where k is a constant.

The minimum velocity of P occurs at $t = 6$

- (i) Show that $k = -7.2$ [4]

- (ii) Find the times at which the particle is at rest. [3]

Initially, the displacement of the particle from a fixed point O is 0.5 m.

- (iii) Find the total distance travelled by the particle between $t = 0$ and $t = 10$ [6]

5 Two particles are moving in the same direction along the same straight line on a smooth horizontal surface.

The particles A and B have masses km kg and m kg respectively, where k is a positive constant.

Initially, the velocities of A and B are 4 m s^{-1} and 2 m s^{-1} respectively, as shown in **Fig. 3** below.



Fig. 3

A collides with B.

After the collision, A moves in the same direction as before, but now with a velocity of 3 m s^{-1}

(i) Find an expression, in terms of k , for the velocity of B after the collision. [4]

(ii) Find an expression, in terms of k and m , for the impulse given to A by B. [3]

(iii) If B is now travelling at twice its original velocity, find the value of k . [2]

6 In this question, take $g = 10 \text{ m s}^{-2}$

Fig. 4 below shows two boxes X and Y each of mass m kg. The boxes are connected by a light inextensible string which passes over a light smooth fixed pulley P. The system lies in a vertical plane.




Fig. 4

X lies on a rough plane inclined at 60° to the horizontal. The coefficient of friction between X and the plane is $\frac{1}{\sqrt{3}}$. Y lies on a smooth plane inclined at 30° to the horizontal.

(i) Draw a diagram showing the external forces acting on the boxes. [2]

The system is released from rest and box X starts to move down the plane.

(ii) Find the acceleration of the system. [8]

(iii) Find, in terms of m , the magnitude of the resultant force exerted by the string on the pulley. [4]

- 7 **Fig. 5** below shows a uniform ladder AB of length 2 m and mass 20 kg.
End A rests on rough horizontal ground.
End B leans against a smooth vertical wall.
The coefficient of friction between the ladder and the ground is 0.3
The ladder is inclined at 25° to the vertical.




Fig. 5

Robert, of mass 70 kg, starts to climb the ladder from A.

- (i) Draw a diagram showing the external forces acting on the ladder. [2]
- (ii) Find how far up the ladder Robert can climb before the ladder starts to slip. [10]

THIS IS THE END OF THE QUESTION PAPER
