



ADVANCED
General Certificate of Education
2015

Mathematics
Assessment Unit F2
assessing
Module FP2: Further Pure Mathematics 2
[AMF21]

TUESDAY 9 JUNE, MORNING

**MARK
SCHEME**

Introduction

The mark scheme normally provides the most popular solution to each question. Other solutions given by candidates are evaluated and credit given as appropriate; these alternative methods are not usually illustrated in the published mark scheme.

The marks awarded for each question are shown in the right-hand column and they are prefixed by the letters **M**, **W** and **MW** as appropriate. The key to the mark scheme is given below:

M indicates marks for correct method.

W indicates marks for working.

MW indicates marks for combined method and working.

The solution to a question gains marks for correct method and marks for an accurate working based on this method. Where the method is not correct no marks can be given.

A later part of a question may require a candidate to use an answer obtained from an earlier part of the same question. A candidate who gets the wrong answer to the earlier part and goes on to the later part is naturally unaware that the wrong data is being used and is actually undertaking the solution of a parallel problem from the point at which the error occurred. If such a candidate continues to apply correct method, then the candidate's individual working must be followed through from the error. If no further errors are made, then the candidate is penalised only for the initial error. Solutions containing two or more working or transcription errors are treated in the same way. This process is usually referred to as "follow-through marking" and allows a candidate to gain credit for that part of a solution which follows a working or transcription error.

Positive marking:

It is our intention to reward candidates for any demonstration of relevant knowledge, skills or understanding. For this reason we adopt a policy of **following through** their answers, that is, having penalised a candidate for an error, we mark the succeeding parts of the question using the candidate's value or answers and award marks accordingly.

Some common examples of this occur in the following cases:

- (a) a numerical error in one entry in a table of values might lead to several answers being incorrect, but these might not be essentially separate errors;
- (b) readings taken from candidates' inaccurate graphs may not agree with the answers expected but might be consistent with the graphs drawn.

When the candidate misreads a question in such a way as to make the question easier only a proportion of the marks will be available (based on the professional judgement of the examining team).

		AVAILABLE MARKS
1	(i) $\begin{aligned} \frac{1}{r^2} - \frac{1}{(r+1)^2} &= \frac{(r+1)^2 - r^2}{r^2(r+1)^2} \\ &= \frac{2r+1}{r^2(r+1)^2} \end{aligned}$	MW1 W1
	(ii) $\begin{aligned} \sum_{r=1}^n \frac{2r+1}{r^2(r+1)^2} &= \sum_{r=1}^n \left[\frac{1}{r^2} - \frac{1}{(r+1)^2} \right] \\ &= \left(1 - \frac{1}{4}\right) + \left(\frac{1}{4} - \frac{1}{9}\right) + \dots + \left(\frac{1}{(n-1)^2} - \frac{1}{n^2}\right) + \left(\frac{1}{n^2} - \frac{1}{(n+1)^2}\right) \\ &= 1 - \frac{1}{(n+1)^2} \end{aligned}$	M1 W1 W1
2	$\tan 2x + \frac{2 \tan 2x}{1 - \tan^2 2x} = 0$ $\tan 2x (1 - \tan^2 2x) + 2 \tan 2x = 0$ $\tan 2x (3 - \tan^2 2x) = 0$ $\tan 2x = 0$ $\therefore 2x = n\pi \quad x = \frac{n\pi}{2}$ or $\tan^2 2x = 3$ $\tan 2x = \pm \sqrt{3}$ $2x = \pm \frac{\pi}{3} + n\pi$ $x = \pm \frac{\pi}{6} + \frac{n\pi}{2}$ Answers can be summarised as $x = \frac{n\pi}{6}$ (not required)	M1 MW1 MW1 W1 MW1 W1 W1 7

			AVAILABLE MARKS	
3	(i)	$\frac{x - 4x^2}{(2-x)^2(3+x^2)} = \frac{A}{2-x} + \frac{B}{(2-x)^2} + \frac{Cx+D}{(3+x^2)}$ $= \frac{A(2-x)(3+x^2) + B(3+x^2) + (Cx+D)(2-x)^2}{(2-x)^2(3+x^2)}$ $x - 4x^2 \equiv A(2-x)(3+x^2) + B(3+x^2) + (Cx+D)(2-x)^2$	M1	
		Let $x = 2$ $-14 = 7B \therefore B = -2$	M1	
		Let $x = 0$ $0 = 6A + 3B + 4D \therefore 6A + 4D = 6$ (a)		
		Coeff. of x^3 $0 = -A + C \therefore A = C$		
		Coeff. of x $1 = -3A + 4C - 4D \therefore A - 4D = 1$ (b)		
		Using (a) and (b) $\Rightarrow A = 1, D = 0$		
		$\therefore C = 1$		
		$\frac{x - 4x^2}{(2-x)^2(3+x^2)} = \frac{1}{2-x} - \frac{2}{(2-x)^2} + \frac{x}{3+x^2}$	W3	
	(ii)	$\int_0^1 \frac{x - 4x^2}{(2-x)^2(3+x^2)} dx = \int_0^1 \frac{1}{2-x} - \frac{2}{(2-x)^2} + \frac{x}{3+x^2} dx$ $= \left[-\ln(2-x) - \frac{2}{2-x} + \frac{1}{2} \ln(3+x^2) \right]_0^1$ $= \left[0 - 2 + \frac{1}{2} \ln 4 \right] - \left[-\ln 2 - 1 + \frac{1}{2} \ln 3 \right]$ $= -1 + \frac{1}{2} \ln \frac{16}{3}$	M1	
			MW3	
			MW1	11
4	(i)	$f(x) = e^{\tan x}$ $f(0) = 1$ $f'(x) = \sec^2 x e^{\tan x}$ $f'(0) = 1$ $f''(x) = \sec^4 x e^{\tan x} + 2 \sec^2 x \tan x e^{\tan x}$ $f''(0) = 1$ $f(x) = f(0) + x f'(0) + \frac{x^2}{2!} f''(0)$ $e^{\tan x} = 1 + x + \frac{x^2}{2}$	MW3	
			M1	
			W1	
	(ii)	$e^{-\tan x} = e^{\tan(-x)} = 1 - x + \frac{x^2}{2}$	MW1	6

5 Let $n = 1$ $3^3 + 2^3 = 35$ is divisible by 7

AVAILABLE MARKS

True for $n = 1$

MW1

Assume true for $n = k$ i.e. $3^{2k+1} + 2^{k+2}$ is divisible by 7

M1

i.e. $3^{2k+1} + 2^{k+2} = 7p$ where p is an integer

Let $n = k + 1$

M1

$$3^{2k+3} + 2^{k+3} = 9 \cdot 3^{2k+1} + 2 \cdot 2^{k+2}$$

$$= 9(7p - 2^{k+2}) + 2 \cdot 2^{k+2} \text{ from assumption}$$

MW1

$$= 63p - 7 \cdot 2^{k+2} = 7(9p - 2^{k+2})$$

\therefore divisible by 7

W1

True for $n = k \Rightarrow$ true for $n = k + 1$

True for $n = 1$

\therefore true for all n

MW1

6

		AVAILABLE MARKS
6	(a) (i) $\frac{2x}{16} + 2y \frac{dy}{dx} = 0$	M1
	$\frac{dy}{dx} = \frac{-x}{16y}$	
	P (4 cos t, sin t) $\frac{dy}{dx} = \frac{-\cos t}{4 \sin t}$	MW1
	$y - \sin t = \frac{-\cos t}{4 \sin t} (x - 4 \cos t)$	MW1
	$4y \sin t - 4 \sin^2 t = -x \cos t + 4 \cos^2 t$	
	$4y \sin t = -x \cos t + 4$	W1
	(ii) Q is point $(0, \frac{1}{\sin t})$ R is point $(\frac{4}{\cos t}, 0)$	MW1 W1
	Midpoint M $(\frac{2}{\cos t}, \frac{1}{2 \sin t})$	MW1
	$x = \frac{2}{\cos t} \Rightarrow \cos t = \frac{2}{x}; y = \frac{1}{2 \sin t} \Rightarrow \sin t = \frac{1}{2y}$	
	$\sin^2 t + \cos^2 t = 1$	M1
	$\therefore \frac{4}{x^2} + \frac{1}{4y^2} = 1 \quad \text{or} \quad \frac{16}{x^2} + \frac{1}{y^2} = 4$	W1
(b)	At intersection	
	$\frac{x^2}{a^2} + \frac{(mx + c)^2}{b^2} = 1$	M1
	$(b^2 + a^2 m^2) x^2 + 2mc a^2 x + a^2 c^2 - a^2 b^2 = 0$	W1
	To be tangent roots equal \therefore discriminant = 0	
	$\therefore (2mc a^2)^2 - 4(b^2 + a^2 m^2)(a^2 c^2 - a^2 b^2) = 0$	M1
	$m^2 c^2 a^2 - b^2 c^2 + b^4 - a^2 c^2 m^2 + a^2 m^2 b^2 = 0$	
	$a^2 m^2 + b^2 = c^2$	W1
		13

			AVAILABLE MARKS
7	Auxiliary Equation	$n^2 + 4n + 13 = 0$	M1
		$n = -2 \pm 3i$	W1
	Complementary Function	$x = e^{-2t}(A \cos 3t + B \sin 3t)$	MW1
	Particular Integral	$x = C \cos 3t + D \sin 3t$	M1
		$\frac{dx}{dt} = -3C \sin 3t + 3D \cos 3t$	MW1
		$\frac{d^2x}{dt^2} = -9C \cos 3t - 9D \sin 3t$	MW1
		$-9C \cos 3t - 9D \sin 3t - 12C \sin 3t + 12D \cos 3t + 13C \cos 3t + 13D \sin 3t = 40 \cos 3t$	
	$12D + 4C = 40$	$-12C + 4D = 0$	
	$D = 3$	$C = 1$	W1 W1
	General Solution	$x = e^{-2t}(A \cos 3t + B \sin 3t) + \cos 3t + 3 \sin 3t$	M1
	$x = 2$ when $t = 0$		
		$\therefore 2 = A + 1 \quad \therefore A = 1$	MW1
		$\frac{dx}{dt} = -2e^{-2t}(A \cos 3t + B \sin 3t) + e^{-2t}(-3A \sin 3t + 3B \cos 3t) - 3 \sin 3t + 9 \cos 3t$	M1
	$\frac{dx}{dt} = 13$ when $t = 0$		
		$\therefore 13 = -2A + 3B + 9 \quad \therefore B = 2$	W1
	Solution	$x = e^{-2t}(\cos 3t + 2 \sin 3t) + \cos 3t + 3 \sin 3t$	W1
			13

				AVAILABLE MARKS
8	(a) (i)	$r = \sqrt{4^2 + 4^2} = 4\sqrt{2}$ $\theta = \frac{\pi}{4}$	MW1 MW1	
	(ii)	$z^5 = [r^5, 5\theta]$	M1	
		$r^5 = 4\sqrt{2}$ $\therefore r = \sqrt{2}$	W1	
		$5\theta = 2n\pi + \frac{\pi}{4}$	$n = 0 \quad \theta = \frac{\pi}{20}$	M1
			$n = 1 \quad \theta = \frac{9\pi}{20}$	
			$n = 2 \quad \theta = \frac{17\pi}{20}$	
			$n = 3 \quad \theta = \frac{25\pi}{20} = \frac{5\pi}{4}$	
			$n = 4 \quad \theta = \frac{33\pi}{20}$	W2
		Solutions	$\sqrt{2} e^{\frac{\pi i}{20}}, \sqrt{2} e^{\frac{9\pi i}{20}}, \sqrt{2} e^{\frac{17\pi i}{20}}, \sqrt{2} e^{\frac{5\pi i}{4}}, \sqrt{2} e^{\frac{33\pi i}{20}}$	W1
	(b)	$\left(\cos \frac{\pi}{8} + i \sin \frac{\pi}{8}\right)^p = e^{\frac{p\pi i}{8}}$		MW1
		$\left(\cos \frac{\pi}{6} - i \sin \frac{\pi}{6}\right)^q = e^{\frac{-q\pi i}{6}}$		MW1
		$-1 = e^{\pi i}$		
		$\frac{\left(\cos \frac{\pi}{8} + i \sin \frac{\pi}{8}\right)^p}{\left(\cos \frac{\pi}{6} - i \sin \frac{\pi}{6}\right)^q} = e^{\frac{p\pi i}{8} + \frac{p\pi i}{6}} = e^{\pi i}$		MW1
		$\frac{p\pi}{8} + \frac{q\pi}{6} = \pi$		W1
		$3p + 4q = 24$		
		p, q , smallest positive integers		
		$\Rightarrow p = 4$ and $q = 3$	W1 W1	14
			Total	75